

1. A simple, safe bimanual technique for subincisional cortex aspiration.
2002 Ophthalmic Surg Lasers.33(4):337-339.
Sakamoto T, Shiraki K, Inoue K, Yanagihara N, Ataka S, Kurita K.
2. Histopathological study of the anterior lens capsule with a double-ring sign.
2004 Graefes Arch Clin Exp Ophthalmol. 242(3):245-249.
Ataka S, Kohno T, Kurita K, Wada S, Takahashi Y, Shiraki K.
3. Quantification and anatomic distribution of choroidal abnormalities in patients with type I neurofibromatosis.
2005 Graefes Arch Clin Exp Ophthalmol. 243(10):980-984.
Nakakura S, Shiraki K, Yasunari T, Hayashi Y, Ataka S, Kohno T.
4. Relation between office intraocular pressure and 24-hour intraocular pressure in patients with primary open-angle glaucoma treated with a combination of topical antiglaucoma eye drops.
2007 J Glaucoma. 16(2):201-204.
Nakakura S, Nomura Y, Ataka S, Shiraki K.
5. Iris prolapse after non-penetrating trabeculectomy with sinusotomy and mitomycin C.
2007 Osaka City Med J. 53(1):49-52.
Takahashi Y, Ataka S, Wada S, Nomura Y, Kohno T, Shiraki K.
6. Multicenter survey with a systematic overview of acute-onset endophthalmitis after transconjunctival microincision vitrectomy surgery.
2010 Am J Ophthalmol. 150(5):716-725
Oshima Y, Kadonosono K, Yamaji H, Inoue M, Yoshida M, Kimura H, Ohji M, Shiraga F, Hamasaki T; Japan Microincision Vitrectomy Surgery Study Group.
7. Dietary glycemic index is inversely associated with the risk of Parkinson's disease: a case-control study in Japan.
2010 Nutrition. 26(5):515-21.
Murakami K, Miyake Y, Sasaki S, Tanaka K, Fukushima W, Kiyohara C, Tsuboi Y, Yamada T, Oeda T, Miki T, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M; Fukuoka Kinki Parkinson's Disease Study Group.

8. Dietary fat intake and risk of Parkinson's disease: a case-control study in Japan.
2010 *J Neurol Sci.* 288(1-2):117-22.
Miyake Y, Sasaki S, Tanaka K, Fukushima W, Kiyohara C, Tsuboi Y, Yamada T, Oeda T, Miki T, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's Disease Study Group.
9. Active and passive smoking and risk of Parkinson's disease.
2010 *Acta Neurol Scand.* 122(6):377-82.
Tanaka K, Miyake Y, Fukushima W, Sasaki S, Kiyohara C, Tsuboi Y, Yamada T, Oeda T, Miki T, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's disease Study Group.
10. Dietary intake of folate, vitamin B6, vitamin B12 and riboflavin and risk of Parkinson's disease: a case-control study in Japan.
2010 *Br J Nutr.* 104(5):757-64.
Murakami K, Miyake Y, Sasaki S, Tanaka K, Fukushima W, Kiyohara C, Tsuboi Y, Yamada T, Oeda T, Miki T, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's Disease Study Group.
11. Case-control study of risk of Parkinson's disease in relation to hypertension, hypercholesterolemia, and diabetes in Japan.
2010 *J Neurol Sci.* 15;293(1-2):82-6.
Miyake Y, Tanaka K, Fukushima W, Sasaki S, Kiyohara C, Tsuboi Y, Yamada T, Oeda T, Miki T, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's Disease Study Group.
12. GST polymorphisms, interaction with smoking and pesticide use, and risk for Parkinson's disease in a Japanese population.
2010 *Parkinsonism Relat Disord.* 16(7):447-52.
Kiyohara C, Miyake Y, Koyanagi M, Fujimoto T, Shirasawa S, Tanaka K, Fukushima W, Sasaki S, Tsuboi Y, Yamada T, Oeda T, Miki T, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's Disease Study Group.
13. LRRK2 Gly2385Arg polymorphism, cigarette smoking, and risk of sporadic Parkinson's disease: a case-control study in Japan.
2010 *J Neurol Sci.* 15;297(1-2):15-8.
Miyake Y, Tsuboi Y, Koyanagi M, Fujimoto T, Shirasawa S, Kiyohara C, Tanaka K, Fukushima W, Sasaki S, Yamada T, Oeda T, Miki T, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's Disease Study Group.

14. Alcohol drinking and risk of Parkinson's disease: a case-control study in Japan.
2010 *BMC Neurol.* 5;10:111.
Fukushima W, Miyake Y, Tanaka K, Sasaki S, Kiyohara C, Tsuboi Y, Yamada T, Oeda T, Miki T, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's Disease Study Group.
15. Lack of association of dairy food, calcium, and vitamin D intake with the risk of Parkinson's disease: a case-control study in Japan.
2010 *Parkinsonism Relat Disord.* 17(2):112-6.
Miyake Y, Tanaka K, Fukushima W, Sasaki S, Kiyohara C, Tsuboi Y, Yamada T, Oeda T, Miki T, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's Disease Study Group.
16. Progression of cataracts following photodynamic therapy combined with intravitreous triamcinolone injection in cases of age-related macular degeneration.
2011 *Osaka City Med J.* 57(2):49-57.
Iwami H, Kohno T, Yamamoto M, Kaida M, Miki N, Ataka S, Shiraki K.
17. Dietary intake of antioxidant vitamins and risk of Parkinson's disease: a case-control study in Japan.
2011 *Eur J Neurol.* 18(1):106-13.
Miyake Y, Fukushima W, Tanaka K, Sasaki S, Kiyohara C, Tsuboi Y, Yamada T, Oeda T, Miki T, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's Disease Study Group.
18. Lack of association of dairy food, calcium, and vitamin D intake with the risk of Parkinson's disease: a case-control study in Japan.
2011 *Parkinsonism Relat Disord.* 17(2):112-6.
Miyake Y, Tanaka K, Fukushima W, Sasaki S, Kiyohara C, Tsuboi Y, Yamada T, Oeda T, Miki T, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's Disease Study Group.
19. APOE and CYP2E1 polymorphisms, alcohol consumption, and Parkinson's disease in a Japanese population.
2011 *J Neural Transm (Vienna).* 118(9):1335-44.
Kiyohara C, Miyake Y, Koyanagi M, Fujimoto T, Shirasawa S, Tanaka K, Fukushima W, Sasaki S, Tsuboi Y, Yamada T, Oeda T, Miki T, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's Disease Study Group.

20. Intake of Japanese and Chinese teas reduces risk of Parkinson's disease.
2011 *Parkinsonism Relat Disord.* 17(6):446-50.
Tanaka K, Miyake Y, Fukushima W, Sasaki S, Kiyohara C, Tsuboi Y, Yamada T, Oeda T, Miki T, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's Disease Study Group.
21. Dietary intake of metals and risk of Parkinson's disease: a case-control study in Japan.
2011 *J Neurol Sci.* 15;306(1-2):98-102.
Miyake Y, Tanaka K, Fukushima W, Sasaki S, Kiyohara C, Tsuboi Y, Yamada T, Oeda T, Miki T, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's Disease Study Group.
22. Occupational risk factors for Parkinson's disease: a case-control study in Japan.
2011 *BMC Neurol.* 7;11:83.
Tanaka K, Miyake Y, Fukushima W, Sasaki S, Kiyohara C, Tsuboi Y, Yamada T, Oeda T, Miki T, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's Disease Study Group.
23. Genetic polymorphisms involved in dopaminergic neurotransmission and risk for Parkinson's disease in a Japanese population.
2011 *BMC Neurol.* 25;11:89.
Kiyohara C, Miyake Y, Koyanagi M, Fujimoto T, Shirasawa S, Tanaka K, Fukushima W, Sasaki S, Tsuboi Y, Yamada T, Oeda T, Shimada H, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's Disease Study Group.
24. Dietary patterns and risk of Parkinson's disease: a case-control study in Japan.
2012 *Eur J Neurol.* 19(5):681-8.
Okubo H, Miyake Y, Sasaki S, Murakami K, Tanaka K, Fukushima W, Kiyohara C, Tsuboi Y, Yamada T, Oeda T, Shimada H, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's Disease Study Group.
25. SNCA polymorphisms, smoking, and sporadic Parkinson's disease in Japanese.
2012 *Parkinsonism Relat Disord.* 18(5):557-61.
Miyake Y, Tanaka K, Fukushima W, Kiyohara C, Sasaki S, Tsuboi Y, Yamada T, Oeda T, Shimada H, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's Disease Study Group.
26. UCHL1 S18Y variant is a risk factor for Parkinson's disease in Japan.
2012 *BMC Neurol.* 28;12:62.
Miyake Y, Tanaka K, Fukushima W, Kiyohara C, Sasaki S, Tsuboi Y, Yamada T, Oeda T, Shimada H, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's Disease Study Group.

27. Lack of association between BST1 polymorphisms and sporadic Parkinson's disease in a Japanese population.
2012 J Neurol Sci. 15;323(1-2):162-6.
Miyake Y, Tanaka K, Fukushima W, Kiyohara C, Sasaki S, Tsuboi Y, Yamada T, Oeda T, Shimada H, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's Disease Study Group.
28. MDR1 C3435T polymorphism and interaction with environmental factors in risk of Parkinson's disease: a case-control study in Japan.
2013 Drug Metab Pharmacokinet. 28(2):138-43.
Kiyohara C, Miyake Y, Koyanagi M, Fujimoto T, Shirasawa S, Tanaka K, Fukushima W, Sasaki S, Tsuboi Y, Yamada T, Oeda T, Shimada H, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M;
Fukuoka Kinki Parkinson's Disease Study Group.
29. Subretinal fluid drainage via original retinal breaks for rhegmatogenous retinal detachment.
2014 Can J Ophthalmol.49(3):256-260 2014
Yamaguchi, Ataka S, Shiraki K.
30. PARK16 polymorphisms, interaction with smoking, and sporadic Parkinson's disease in Japan.
2016 J Neurol Sci. 15;362:47-52.
Miyake Y, Tanaka K, Fukushima W, Kiyohara C, Sasaki S, Tsuboi Y, Oeda T, Shimada H, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M, Nakamura Y;
Fukuoka Kinki Parkinson's Disease Study Group.
31. Vitamin D receptor gene polymorphisms, smoking, and risk of sporadic Parkinson's disease in Japan.
2017 Neurosci Lett. 16;643:97-102.
Tanaka K, Miyake Y, Fukushima W, Kiyohara C, Sasaki S, Tsuboi Y, Oeda T, Shimada H, Kawamura N, Sakae N, Fukuyama H, Hirota Y, Nagai M, Nakamura Y;
Fukuoka Kinki Parkinson's Disease Study Group.